



Leaders in Eye Care

[www.eyecenteroftexas.com](http://www.eyecenteroftexas.com)

6565 West Loop South  
Suite 650  
Bellaire, Texas 77401  
Ph. 713.797.1010  
Fax.713.357.7276

4415 Crenshaw  
Pasadena, Texas 77504  
Ph. 281.977.8800  
Fax.281.977.8877

15400 Southwest Frwy  
Suite 301  
Sugar Land, Texas 77478  
Ph. 281.277.1010  
Fax.281.277.4504

21700 Kingsland Blvd.  
Suite 102  
Katy Medical Complex  
Professional Building 1  
Katy, TX 77450  
Ph. 713.797.1010  
Fax.713.357.7276

450 Medical Center Blvd.  
Suite 305  
Webster, Texas 77598  
Ph. 281.332.1397  
Fax 281.338.1215

11914 Astoria Blvd.  
Suite 325  
Memorial SE Prof. Bldg.  
Houston, Texas 77089  
Ph. 281.484.2030  
Fax 281.484.6549

Edward C. Wade, M. D.  
Ting Fang-Suarez, M. D.  
Mark L. Mayo, M. D.  
Gurpreet D. Singh, M. D.

Christopher D. Allee, O. D.  
Jill Autry, O. D.  
Randall N. Reichle, O. D.  
Amanda Bachman, O. D.

## Corneal Transplant

### What is a corneal transplant?

The cornea is normally a clear layer of tissue covering the front of the eye, similar to a watch crystal. Its purpose is to refract or bend light rays as they enter the eye, allowing them to focus on the retina. In cases where the cornea has become clouded as a result of disease, swelling, scarring, infection, chemical burns, a corneal transplant (also called a keratoplasty) is sometimes necessary to restore functional vision. For this procedure, the surgeon carefully removes the damaged central corneal tissue and replaces it with a precisely shaped replica of donor tissue.

### What are the chances of a successful surgery?

The success rate of corneal transplant is about 85%. However, factors such as glaucoma, macular degeneration, or optic nerve disease may affect the final visual result even if the transplant is successful.

The procedure usually takes approximately one hour and is performed with local anesthesia on an outpatient basis. Drops are prescribed before and after surgery to prevent infection and rejection of the donor cornea. After the surgery, a plastic shield should be worn at night for one week to prevent rubbing, bumping, or scratching the eye.

During the postoperative period, a doctor monitors the cornea's healing with special computer mapping called corneal topography. This allows the doctor to evaluate the shape of the new cornea and remove sutures as needed to control astigmatism.

### What type of vision can you expect after surgery?

In most cases, vision returns very gradually. The healing process may vary greatly from one individual to the next. Some may enjoy improved vision within a few months after surgery, for others, it may take up to a year. In most cases, the use of a special rigid contact lens is required for optimal vision.