



6565 West Loop South
Suite 650
Bellaire, Texas 77401
Ph. 713.395.1515
Fax.713.357.7278

4415 Crenshaw Rd.
Pasadena, Texas 77504
Ph. 281.977.8800
Fax.281.977.8877

15400 Southwest Frwy
Suite 301
Sugar Land, Texas 77478
Ph. 281.277.1010
Fax.281.277.4504

450 Medical Center Blvd.
Suite 305
Webster, Texas 77598
Ph. 281.332.1397
Fax 281.338.1215

21700 Kingsland Blvd.
Suite 102
Katy Medical Complex
Professional Building 1
Katy, TX 77450
Ph. 713.797.1010
Fax.713.797.6200

Edward C. Wade, M. D. Christopher D. Allee, O. D.
Ting Fang-Suarez, M. D. Jill Autry, O. D.
Mark L. Mayo, M. D. Amanda Bachman, O. D.
Randall N. Reichle, O. D. Julie Ngo, O. D.

How to Use Latisse

LATISSE® solution is intended for **use on the skin of the upper eyelid margins at the base of the eyelashes**. Refer to the illustration below. **DO NOT APPLY** to the lower eyelid. If you are using LUMIGAN® or other products in the same class for elevated intraocular pressure (IOP), or if you have a history of abnormal IOP, you should only use LATISSE® under the close supervision of your physician.



LATISSE® use may cause darkening of the eyelid skin which may be reversible. LATISSE® use may also cause increased brown pigmentation of the colored part of the eye which is likely to be permanent.

It is possible for hair growth to occur in other areas of your skin that LATISSE® frequently touches. Any excess solution outside the upper eyelid margin should be blotted with a tissue or other absorbent material to reduce the chance of this from happening. It is also possible for a difference in eyelash length, thickness, fullness, pigmentation, number of eyelash hairs, and/or direction of eyelash growth to occur between eyes. These differences, should they occur, will usually go away if you stop using LATISSE®.

Other Important Safety Information

DO NOT APPLY in your eye or to the lower lid. **ONLY** use the sterile applicators supplied with LATISSE® to apply the product.

Don't allow the tip of the bottle or applicator to contact surrounding structures, fingers, or any other unintended surface in order to avoid contamination by common bacteria known to cause infections.

Contact lenses should be removed prior to application of LATISSE® and may be reinserted 15 minutes following its administration.