

Edward C. Wade, M. D Ting Fang-Suarez, M. D. Mark L. Mayo, M. D. Randall N. Reichle, O. D Christopher D. Allee, O. D. Jill Autry, O. D. Amanda Bachman, O. D. Julie Ngo, O. D.

www.eyecenteroftexas.com

6565 West Loop South Suite 650 Bellaire, Texas 77401 Ph. 713.395.1515 Fax.713.357.7278

4415 Crenshaw Rd. Pasadena, Texas 77504 Ph. 281.977.8800 Fax.281.977.8877

15400 Southwest Frwy Suite 301 Sugar Land, Texas 77478 Ph. 281.277.1010 Fax.281.277.4504

450 Medical Center Blvd. Suite 305 Webster, Texas 77598 Ph. 281.332.1397 Fax 281.338.1215

21700 Kingsland Blvd. Suite 102 Katy Medical Complex Professional Building 1 Katy, TX 77450 Ph. 713.797.1010 Fax.713.797.6200

Lasik Pre-Op Instructions

Patient Name:		
Surgeon:		
Date of Surgery:	Arrival Time:	

- 1. You <u>must</u> have a responsible adult with you the day of surgery. You <u>must</u> have someone stay with you the night after your surgery.
- You <u>must</u> see one of our doctors the day after your surgery. Take your regular morning medications (if any) with breakfast with the following exceptions: <u>No</u> aspirin 5 days prior to surgery; no Accutane four (4) weeks prior to surgery. Restart Accutane four (4) weeks <u>after</u> surgery. <u>If you are taking Amiodarone</u>, you are **not** a <u>LASIK</u> candidate. Please check with your medical doctor before making any changes in your medications.
- 3. Remove eye makeup thoroughly the night prior to surgery. Do not wear eye makeup, earrings, nail polish, perfume, cologne, after-shave, hair spray, scented deodorant or anything that has a <u>fragrance</u>, as <u>this will affect the laser beam's effect on your eye</u>! Leave all jewelry and valuables at home. Do not wear your hair in a ponytail or braid.
- 4. Do not wear soft contact lenses in the operative eye(s) for 2 weeks prior to evaluation and surgery.
- 5. Do not wear hard or gas permeable contact lenses in the operative eye(s) for 3 weeks prior to evaluation and surgery. You can substitute a soft contact lens until 2 weeks before surgery.
- 6. It is recommended that you eat a light breakfast before coming for your surgery. Be sure to have your prescription filled <u>before</u> the surgery and <u>bring</u> the medication with you when you report to the laser center. <u>Please use the</u> <u>ZYMAXID, if prescribed, 3 days prior to surgery</u>. Use Restasis, if prescribed by your doctor, 3 days prior to surgery.
- 7. Plan on being at the laser center for <u>about 2 hours.</u>
- 8. Report to: Eye Center Of Texas

6565 West Loop South, Suite 650,

Bellaire, Texas 77401

(Located between Bellaire and Bissonnet on the inside 610 Loop Service Road).

If you have any questions about these instructions, please call

EYE CENTER OF TEXAS

713-395-1515, 713-357-7229, 713-357-7230 or call your optometrist.